



**UNIVERSITI PUTRA MALAYSIA**

**PHASE BEHAVIOUR PHYSICAL PROPERTIES OF BATYL  
ALCOHOL**

**LIM CHAI SENG.**

**FSAS 2004 9**

**PHASE BEHAVIOUR AND PHYSICAL PROPERTIES  
OF BATYL ALCOHOL**

**By**

**LIM CHAI SENG**

**Thesis Submitted to the School of Graduate Studies,  
Universiti Putra Malaysia, in Fulfilment of  
the Requirements for the Degree of Master**

**September 2004**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirements for the degree of Master of Science

**PHASE BEHAVIOUR AND PHYSICAL PROPERTIES  
OF BATYL ALCOHOL**

**By**

**LIM CHAI SENG**

**June 2004**

**Chairman: Professor Anuar Kassim, Ph.D**

**Faculty: Science and Environmental Studies**

Batyl alcohol (monoalkylether glycerine) acts as an emulsifier when used in emulsion products, such as cosmetics creams, lotions and ointments. It is chemically stable, highly purified and safe. Although batyl alcohol is widely used in cosmetic products, little is known about its phase behaviour and rheological properties in ternary systems. Its phase behaviour in ternary systems was therefore studied at 80°C, and observed under a polarising microscope.

In the ternary systems, lamellar and hexagonal areas were found in 90/10 and 100/0 mixtures of batyl alcohol (BA) and isopropyl myristate (IPM) and the percentage of water were from 9% - 44% w/w, respectively. They were identified by their patterns of maltese crosses and fan structures, respectively.

Both of them exhibited a viscoelastic network. Further addition of either medium chain triglycerides (MCT) or propylene glycol (PG) to the system resulted in isotropic and two-phase areas being formed instead of a liquid crystalline structure.

Using a combination of BA, IPM, MCT and water, an emulsion was developed. A ratio of 1:1 MCT:water was emulsified with 5% and 10% BA. A weak viscoelastic network was formed with 5% BA but without stearic acid (Emulsion S1). With stearic acid (Emulsion S2) a strong viscoelastic network was formed with the acid acting as a co-emulsifier. No network structure was found in the emulsion containing 10% BA (Emulsion S4) as it was stabilized by the right percentage of BA. The emulsifiers adsorbed at the MCT/water interface, reducing the droplet size and increasing the viscosity, while the rest formed liquid crystalline lamellar in the continuous phase to physically trap the droplets in the network. The conductivity measured revealed that oil-in-water emulsion can conduct electric and it also shows the oil droplets encapsulated by the emulsifier in S2.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**KELAKUAN FASA DAN CIRI-CIRI FIZIKAL  
BAGI BATIL ALKOHOL**

**Oleh**

**LIM CHAI SENG**

**Jun 2004**

**Pengerusi:           Profesor Anuar Kassim, Ph.D**

**Fakulti:             Sains dan Pengajian Alam Sekitar**

Batil alkohol (monoalkileter gliserin), apabila digunakan dalam barangan emulsi seperti krim kosmetik, losyen dan salap, ia akan bertindak sebagai pengemulsi. Sebatian ini adalah stabil dari segi kimia, sangat tulen dan selamat digunakan. Walaupun batil alkohol digunakan secara meluas dalam barangan kosmetik, pengetahuan terhadap kelakuan fasa dan sifat reologi dalam sistem tiga fasa agak kurang. Kelakuan fasa dalam sistem tiga fasa telah dikaji pada 80°C serta dilihat melalui cahaya berpengutub dan disahkan dengan mikroskop berpengutub.

Dalam sistem tiga fasa, fasa lamelar dan heksagonal terdapat pada kombinasi 90/10 dan 100/0 batil alcohol (BA)/ isopropil miristat (IPM) dan peratus air adalah dari 9% - 44% w/w masing-masing dan mereka telah

diidentifikasi sebagai corak palang maltese dan taring masing-masing. Keduanya menunjukkan jalinan hablur cecair yang bersifat viskoelastik. Isotropik dan kawasan dua fasa telah terbentuk dengan penambahan trigliserida rantai sederhana (MCT) dan propilena glikol (PG) kepada sistem ini.

Dengan campuran BA, IPM, MCT dan air, sistem emulsi telah dimajukan. Nisbah 1:1 bagi MCT dan air telah diemulsi dengan 5% dan 10% BA. Terdapat satu viskoelastik yang lemah telah terbentuk dalam sampel S1 dengan 5% BA tanpa asid stearik tetapi dengan kehadiran asid stearik dalam sampel S2, satu viskoelastik yang kuat telah terbentuk. Asid stearik bertindak sebagai pengemulsi bersama. Emulsi yang mengandungi 10% BA dalam S4 tidak terdapat sebarang viskoelastik. Sistem distabilkan dengan menggunakan peratusan BA yang betul. Pengemulsi menjerap pada antara muka MCT/air, mengurangkan saiz titisan, meningkatkan kelikatan, semasa yang lain membentuk lamelar hablur cecair dalam fasa terusan sehingga memerangkap titisan secara fizikal. Pengukuran kekonduksian menunjukkan emulsi minyak dalam air boleh mengkonduksi elektrik dan ia juga menunjukkan titisan minyak diperangkap oleh pengemulsi dalam sistem S2.

## **ACKNOWLEDGEMENTS**

I would like to express my sincere appreciation to my supervisor, Prof. Dr. Anuar Kassim, for his invaluable advice, guidance and suggestions throughout this project. My deepest gratitude is also due my co-supervisor, Dr. Zahariah Ismail, for her ideas, unfailing help, constructive criticism and support. This is, of course, not to forget Mr. Halim from AOTC and Mr. Chang Kwong Choong from MPOB for all their help.

I would like to thank AOTC for allowing me to use its facilities. Last but not the least, I would like to thank my family, beloved friends and fellow laboratory mates for their support and encouragement.

I certify that an Examination Committee met on 9<sup>th</sup> June 2004 to conduct the final examination of Lim Chai Seng on his Master of Science thesis entitled "Phase Behaviour and Physical Properties of Batyl Alcohol" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Mohd. Zaizi Desa, Ph.D.**

Associate Professor  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Chairman)

**Anuar Kassim, Ph.D.**

Professor  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)

**Zahariah Ismail, Ph.D.**

Advanced Oleochemicals Technology Centre  
Malaysian Palm Oil Board  
(Member)

**Dzulkefly Kuang, Ph.D.**

Professor  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)



---

**GULAM/RUSUL RAHMAT ALI, Ph.D.**  
Professor/Deputy Dean  
School of Graduates Studies  
Universiti Putra Malaysia

Date: 08 SEP 2004



This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Anuar Kassim, Ph.D.**

Professor  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Chairman)

**Zahariah Ismail, Ph.D.**

Advanced Oleochemical Technology Centre  
Malaysian Palm Oil Board  
(Member)

**Dzulkefly Kuang, Ph.D.**

Professor  
Faculty of Science and Environmental Studies  
Universiti Putra Malaysia  
(Member)




---

**AINI IDERIS, Ph.D.**  
Professor/ Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 10 SEP 2004

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or any other institutions.



LIM CHAI SENG

Date: 20 AUG 2004

## TABLE OF CONTENTS

	<b>Page</b>
ABSTRACT	ii
ABSTRAK	iv
ACKNOWLEDGEMENTS	vi
APPROVAL	vii
DECLARATION	ix
LIST OF FIGURES	xiii
LIST OF TABLES	xvi
LIST OF ABBREVIATIONS	xviii
 <b>CHAPTER</b>	
<b>1 INTRODUCTION</b>	
1.1 Liquid Crystal	1
1.1.1 Lyotropic Liquid Crystal	3
1.1.2 Classification of Lyotropic Liquid Crystal	4
1.1.3 Lamellar Phase	5
1.1.4 Hexagonal Phase	7
1.2 Emulsions	9
1.2.1 Emulsifiers	10
1.2.2 Functions of Emulsifiers	10
1.2.3 Emollients	12
1.2.4 Hydrophile-Lipophile Balance	13
1.2.5 Emulsification	16
1.3 Rheology	17
1.3.1 Classification of Materials	19
1.3.2 Viscosity	21
1.3.3 Thixotropy	22
1.4 Conductivity	24
1.5 Objectives of the Study	25
 <b>2 LITERATURE REVIEW</b>	
2.1 Batyl Alcohol	26
2.2 Lamellar Liquid Crystal and Rheological Properties in Emulsion Systems	27
2.3 Emulsifying Properties Using the Conductivity Technique	31
 <b>3 MATERIALS AND METHODS</b>	
3.1 Materials	33
3.1.1 Monostearyl Glycerine Ether	33
3.1.2 Isopropyl Myristate	34
3.1.3 Triethanolamine	35



3.1.4	Glycerol Tricapryllate/ Tricaprate	36
3.1.5	Steric Acid	38
3.1.6	Propylene Glycol	39
3.2	Methods	40
3.2.1	Constructing the Ternary Phase Diagrams	40
3.2.1.1	Preparing the Ternary of BA/IPM/H <sub>2</sub> O System	40
3.2.1.2	Preparing the Ternary BA/IPM (90/10)/H <sub>2</sub> O/MCT System	41
3.2.1.3	Preparing the Ternary BA/IPM (90/10)/H <sub>2</sub> O/PG System	41
3.2.1.4	Preparing the Ternary BA/IPM (100/0)/H <sub>2</sub> O/MCT System	41
3.2.1.5	Preparing the Ternary BA/IPM (100/0)/H <sub>2</sub> O/PG System	42
3.2.2	Rheology Measurements	42
3.2.2.1	Sample Preparation for Thixotropy and Viscosity Measurement	42
3.2.2.1.1	Ternary BA/IPM/ H <sub>2</sub> O System	42
3.2.2.1.2	Ternary BA/IPM (90/10)/H <sub>2</sub> O/MCT System	43
3.2.2.1.3	Ternary BA/IPM (90/10)/H <sub>2</sub> O/PG System	43
3.2.2.1.4	Ternary BA/IPM (100/0)/H <sub>2</sub> O/MCT System	44
3.2.2.1.5	Ternary BA/IPM (100/0)/H <sub>2</sub> O/PG System	45
3.2.2.2	Thixotropy Measurements for Phase Behaviour and Emulsion System	45
3.2.2.3	Viscosity Measurements for Phase Behaviour and Emulsion System	46
3.2.3	Optical Microscopy Measurements	46
3.2.4	Emulsion Preparation	47
3.2.5	Conductivity Measurements	48
3.2.6	Emulsion Stability Measurements	48
4	RESULTS AND DISCUSSION	
4.1	Phase Behaviour of the ternary BA/IPM/H <sub>2</sub> O System	49
4.2	Phase Behaviour of the ternary BA/IPM (90/10)/H <sub>2</sub> O/MCT System	55
4.3	Phase Behaviour of the ternary BA/IPM (90/10)/H <sub>2</sub> O/PG System	57
4.4	Phase Behaviour of the ternary BA/IPM	



	(100/0)/H <sub>2</sub> O/MCT System	59
4.5	Phase Behaviour of the ternary BA/IPM (100/0)/H <sub>2</sub> O/PG System	61
4.6	Rheology	64
4.6.1	Thixotropic Behaviour of the BA/IPM/H <sub>2</sub> O System	64
4.6.2	Thixotropic Behaviour of the BA/IPM (90/10)/H <sub>2</sub> O/MCT System	67
4.6.3	Thixotropic Behaviour of the BA/IPM (90/10)/H <sub>2</sub> O/PG System	69
4.6.4	Thixotropic Behaviour of the BA/IPM (100/0)/H <sub>2</sub> O/MCT System	70
4.6.5	Thixotropic Behaviour of the BA/IPM (100/0)/H <sub>2</sub> O/PG System	72
4.7	Viscosity and Yield Measurements	74
4.7.1	Phase Diagram of the ternary BA/IPM/ H <sub>2</sub> O System	74
4.7.2	Phase Diagram of the ternary BA/ IPM (90/10)/H <sub>2</sub> O/MCT System	76
4.7.3	Phase Diagram of the ternary BA/ IPM (90/10)/H <sub>2</sub> O/PG System	77
4.7.4	Phase Diagram of the ternary BA/ IPM (100/0)/H <sub>2</sub> O/MCT System	78
4.7.5	Phase Diagram of the ternary BA/ IPM (100/0)/H <sub>2</sub> O/PG System	79
4.8	Emulsion Using Batyl Alcohol	80
4.8.1	Emulsion Droplet Stability and Viscosity under the Microscope	80
4.8.2	Thixotropic Behaviour of the Emulsions	86
4.9	Conductivity Measurements	89
5	CONCLUSION AND FUTURE WORK	93
	REFERENCES	96
	APPENDICES	103
	BIODATA OF THE AUTHOR	136



## LIST OF FIGURES

Figure		Page
1.1	lamellar liquid crystal (a) Schematic structure, and (b) Typical pattern in it i) Maltese crosses ii) oily streak and iii) striation.	6
1.2	(a) Schematic structure of normal ( $H_1$ ) and reversed ( $H_2$ ) hexagonal and (b) Typical pattern of hexagonal liquid crystal i) $H_1$ and ii) $H_2$ .	8
1.3	Material under shear stress.	19
1.4	Flow curve for thixotropic behaviour.	23
3.1	Structure of Batyl alcohol 100.	33
3.2	Structure of isopropyl myristate.	34
3.3	Structure of triethanolamine.	35
3.4	Structure of glycerol tricaprylate/ tricaprates.	37
3.5	Structure of stearic acid.	38
3.6	Structure of propylene glycol.	39
4.1	Phase diagram of the BA/IPM/ $H_2O$ system at 80°C.	51
4.2	Optical pattern of 100/0 BA/IPM in 30% water showing the fan texture of the hexagonal phase. Viewed under crossed-polarized light @ 200X magnification.	52
4.3	Optical pattern of 90/10 BA/IPM in 30% water showing Maltese crosses of the lamellar phase. Viewed under crossed-polarized light @ 200X magnification.	52
4.4	Optical pattern of 90/10 BA/IPM showing the focal conic fan like and pseudo-isotropic texture (in 15% water) of the isotropic phase. Viewed under crossed-polarized light @ 200X magnification.	54
4.5	Phase diagram of the BA/IPM (90/10)/ $H_2O$ /MCT system at 80°C.	56



4.6	Phase diagram of the BA/IPM (90/10)/H <sub>2</sub> O/PG system at 80°C.	58
4.7	Phase diagram of the BA/IPM (100/0)/H <sub>2</sub> O/MCT system at 80°C.	60
4.8	Phase diagram of the BA/IPM (100/0)/H <sub>2</sub> O/PG system at 80°C.	63
4.9	Thixotropic behaviour of the hexagonal phase of 90/10 BA/IPM at 80°C in a) 30% water w/w, and b) 35% water w/w.	66
4.10	Thixotropic behaviour of the lamellar phase of 100/0 BA/IPM at 80°C in a) 30% water w/w, and b) 35% water w/w.	66
4.11	Thixotropic behaviour of the focal-conic fan like and pseudo-isotropic area of 90/10 BA/IPM at 80°C in a) 15% water w/w, and b) 10% water w/w.	67
4.12	Thixotropic behaviour of the isotropic area of 90/10 BA/IPM (90/10)/H <sub>2</sub> O at 80°C in a) 10% MCT w/w, and b) 30% MCT w/w.	68
4.13	Thixotropic behaviour of the two-phase area of 30/70 BA/IPM (90/ 10)/H <sub>2</sub> O at 80°C in a) 25% MCT w/w, 50/50 BA/IPM (90/ 10)/H <sub>2</sub> O, in b) 45% MCT w/w.	68
4.14	Thixotropic behaviour of the isotropic area of 90/10 BA/IPM (90/10)/H <sub>2</sub> O at 80°C in a) 10% PG w/w, and b) 30% PG w/w.	69
4.15	Thixotropic behaviour of the two-phase area of 30/70 BA/IPM (90/10)/H <sub>2</sub> O at 80°C in a) 25% PG w/w, 50/50 BA/IPM (90/10)/H <sub>2</sub> O in b) 45% PG w/w.	70
4.16	Thixotropic behaviour of the isotropic area of 90/10 BA/IPM (100/0)/H <sub>2</sub> O at 80°C in a) 10% MCT w/w, and b) 30% MCT w/w.	71
4.17	Thixotropic behaviour of the two-phase area of 30/70 BA/IPM (100/0)/H <sub>2</sub> O at 80°C in a) 25% MCT w/w, 50/50 BA/IPM (100/0)/H <sub>2</sub> O in b) 45% MCT w/w.	71



4.18	Thixotropic behaviour of the isotropic area of 90/10 BA/IPM (100/0)/H <sub>2</sub> O at 80°C in a) 10% PG w/w, and b) 30% PG w/w.	72
4.19	Thixotropic behaviour of the two-phase area of 30/70 BA/IPM (100/0)/H <sub>2</sub> O at 80°C in a) 25% PG w/w, 50/50 BA/IPM (100/0)/H <sub>2</sub> O in b) 45% PG w/w.	73
4.20	Structure of emulsion S1 at room temperature. Viewed through a light microscope at 50x magnification.	83
4.21	Structure of emulsion S2 at room temperature. Viewed through a light microscope at 50x magnification.	83
4.22	Structure of emulsion S3 at room temperature. Viewed through a light microscope at 50x magnification.	84
4.23	Structure of emulsion S4 at room temperature. Viewed through a light microscope at 50x magnification.	84
4.24	Thixotropic behaviour of emulsion S1.	87
4.25	Thixotropic behaviour of emulsion S2.	87
4.26	Thixotropic behaviour of emulsion S3.	88
4.27	Thixotropic behaviour of emulsion S4.	88
4.28	Conductivity curve of emulsion S1.	91
4.29	Conductivity curve of emulsion S2.	91
4.30	Conductivity curve of emulsion S3.	92
4.31	Conductivity curve of emulsion S4.	92



## LIST OF TABLES

Figure		Page
1.1	HLB by Dispersibility.	15
3.1	Specifications of Batyl alcohol 100.	33
3.2	Specifications of isopropyl myristate.	34
3.3	Specifications of triethanolamine.	35
3.4	Specifications of glycerol tricaprylate/tricaprate	36
3.5	Specifications of stearic acid.	37
3.6	Specifications of propylene glycol.	38
3.7	Emulsion systems using 5% and 10% Batyl alcohol.	43
4.1	Summary of the phase transition of the ternary BA/IPM/H <sub>2</sub> O system at 80°C.	48
4.2	Viscosity and yield value of the ternary BA/IPM/H <sub>2</sub> O system at 80°C and various H <sub>2</sub> O contents (Agitated at 60rpm in a Bingham rheometer).	68
4.3	Viscosity and yield value of the ternary BA/IPM (90/10)/H <sub>2</sub> O/MCT system at 80°C and various MCT contents (Agitated at 60rpm in a Bingham rheometer).	69
4.4	Viscosity and yield value of the ternary BA/IPM (90/10)/H <sub>2</sub> O/PG system at 80°C and various PG contents (Agitated at 60rpm in a Bingham rheometer).	70
4.5	Viscosity and yield value of the ternary BA/IPM (100/0)/H <sub>2</sub> O/MCT system at 80°C and various MCT contents (Agitated at 60rpm in a Bingham rheometer).	71
4.6	Viscosity and yield value of the ternary BA/IPM (100/0)/H <sub>2</sub> O/PG system at 80°C and various PG contents (Agitated at 60 rpm in a Bingham rheometer).	72
4.7	Stability of 50% MCT w/w (emulsified with 5% and 10% Batyl alcohol) after 3 months' storage.	73



4.8	Apparent viscosities at 25°C for a shear rate of 230s <sup>-1</sup> for emulsions S1-S4.	77
-----	--	----



## **LIST OF ABBREVIATIONS**

**BA = Batyl Alcohol**

**IPM = Isopropyl Myristate**

**MCT = Medium Chain Triglyceride**

**PG = Propylene glycol**

**TEA = Triethanolamine**

**SA = Stearic Acid**

## CHAPTER 1

### INTRODUCTION

#### 1.1 Liquid Crystal

Otto Lehmann, a physicist, first coined the term *liquid crystal* to describe materials which share many of the properties of both the liquid and solid states. The first liquid crystalline material discovered was an organic substance related to cholesterol - cholesterol benzoate. It was actually first observed around 1850 by Heintz, a chemist working on natural fats, but only reported in 1888 by Friedrich Reinitzer, an Austrian botanist.

The meaning of "liquid crystal" has been enormously extended in the last two decades to encompass both the chemical and structural concepts. In general, it is accepted that liquid crystals represent a higher state of order than ordinary (isotropic) liquids. However, the delimitation from crystalline solid is less clear. Liquid crystals possess greater intermolecular and intramolecular mobility than classical solids, with many types having a lower degree of order than crystalline solids. However, there are several types of highly ordered smectic

liquid crystals, which possess a three-dimensional order, and are therefore designated "crystal" phase types by Mori *et al.* (1997).

There are three commonly recognized states of matter - solid, liquid and gas. Solid may be either crystalline or amorphous. Crystalline solids have a regular arrangement of their molecules over a large distance compared to their molecular dimensions, or a long-range order. When a crystalline solid is heated, it transforms into an isotropic liquid at its melting point. The isotropic liquid does not have any long-range order. Conversely, on cooling, the isotropic liquid is transformed into a crystalline solid (Ekwall, 1974).

For years after their discovery, liquid crystals remained a scientific curiosity. They were studied by scientists, who thought they had learnt everything about them by the end of the Second World War. The past decade though has witnessed new impetus in liquid crystal research with specific applications in such diverse areas as medicine, biology, chemistry, physics, space science, mathematics and engineering (Westerman, 1993).

Liquid crystal is a material that can transform from the solid to liquid state and *vice versa* with an intermediate phase in between – the mesomorphase. Thus, liquid crystal is similar to a liquid–free flow due to the absence of positional order. On the other hand, liquid crystal also resembles a crystalline solid because it maintains some orientation. Basically, liquid crystal is divided into *lyotropic* and

*thermotropic*. A lyotropic liquid crystal is formed when a surfactant is mixed with a solvent whereas a thermotropic liquid crystal is formed over a certain temperature range (Ekwall, 1974).

### **1.1.1 Lyotropic Liquid Crystal**

The word “lyotropic” means “solvent-induced”, which starts with molecules that are amphiphilic. Amphiphilic molecules are composed of two different parts - a nonpolar, or hydrophobic, hydrocarbon tail insoluble in water, and a polar or hydrophilic head soluble in water. Some amphiphilic molecules that form lyotropic liquid crystals under controlled conditions are soaps, bile salts and phospholipids (Collings, 1990). When these compounds are dissolved in water, they can form spherical aggregates such as micelles, or vesicles, or cylindrical structures such as a bilayer. These structures float freely in the water but retain their orientation and positional order.

It has long been recognized that liquid crystalline phases form a water miscible surfactant, which, when dissolved in water above a well-defined concentration, form aggregates (micelles). As the surfactant concentration increases, the physical characteristics of the solution will change the nature of the aggregated solute. These aggregates are the building blocks of the liquid crystal phases that occur at higher concentration.

### **1.1.2 Classification of Lyotropic Liquid Crystal**


There are six classes of lyotropic liquid crystal - lamellar, hexagonal, cubic, nematic, gel and intermediate phases. All of these have been recognized for many years except the intermediate phase. However, for simplicity, only the two major types are discussed - lamellar and hexagonal. Furthermore, different lamellar liquid crystals are formed from different combinations of interlayer spacings.

### 1.1.3 Lamellar Phase


The most common lyotropic liquid crystal form in a surfactant system is the lamellar phase (Figure 1.1), also known as the neat phase. The lamellar structure is the most extensive liquid crystal phase studied as it has only a simple one-dimensional order. In this phase, the surfactant molecules are arranged in bilayers, separated by water layers.

The surfactant in the bilayer is arranged such that the hydrophobic groups of the surfactant molecules are located in the center of the bilayer. The hydrophilic groups are therefore attached to the solvent layer. Viewed under a polarising microscope, the liquid crystal has three optical patterns – Maltese crosses, oily streaks and striation (Figure 1.1b). The thickness of the bilayer structure is dependent on the water content - increasing the water content will increase the area per polar head group of the occupied interface (Ekwall, 1975; Small, 1988).





(a)



(i)

(ii)

(iii)

(b)

Figure 1.1: Lamellar liquid crystal (a) Schematic structure, and (b) Typical patterns in it: i) Maltese crosses, ii) oily streaks, and iii) striation.